

Abstract Submitted
for the DNP20 Meeting of
The American Physical Society

Imaging of single Ba atoms and Ba⁺ ions in solid xenon for barium tagging in next-generation ¹³⁶Xe double beta decay experiments¹ DAVID FAIRBANK, JAMES TODD, ALEC IVERSON, TREY WAGER, WILLIAM FAIRBANK, Colorado State University, NEXO COLLABORATION COLLABORATION — The identification, or “tagging” of the barium-136 daughter atom that results from double beta decay of xenon-136 provides a promising technique for elimination of all backgrounds except 2-neutrino double beta decay in future generations of ¹³⁶Xe neutrinoless double beta decay experiments. The Ba tagging scheme being developed utilizes a cryogenic probe to trap the ¹³⁶Ba daughter atom in solid xenon and extract it from a liquid xenon time projection chamber, such as the nEXO design concept. The barium atom is then tagged via fluorescence imaging in the solid xenon matrix. The status of our efforts to image and count single Barium atoms and Ba⁺ ions in the solid xenon matrix will be presented. An important feature of the method is that any residual Ba atoms on the probe surface do not create an observable signal, only those that are captured in the solid xenon.

¹Support provided by NSF Grant number 1649324

William Fairbank
Colorado State University

Date submitted: 26 Jun 2020

Electronic form version 1.4