

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Application of GEM Code for Experimentally-Realistic Tokamak Cases¹ G. REWOLDT, W.M. TANG, Princeton Plasma Physics Lab., Y. CHEN, S.E. PARKER, U. Colorado Boulder CIPS — The GEM code is a gyrokinetic electromagnetic nonlinear particle-in-cell simulation code[1]. It has recently been extended to be radially global[2], using the Miller toroidal MHD equilibrium[3]. In recent work, the GEM code has been interfaced with the TRANSP experimental data system[4]. In particular, the GEM code calculations can now include experimentally-derived TRANSP density and temperature profiles for the electron and background ion species, as well as for an impurity species and a hot beam ion species, with trapped electrons and electron collisions, and including self-generated and externally-driven flow. Preliminary results will be presented for tokamaks such as NSTX and DIII-D.

[1] Y. Chen and S. Parker, J. Comput. Phys. **189**, 462 (2003).

[2] Y. Chen and S. Parker, J. Comput. Phys., in press (2006).

[3] R.L. Miller, *et al.*, Phys. Plasmas **5**, 973 (1998); R.E. Waltz, *et al.*, Phys. Plasmas **6**, 4265 (1999).

[4] R. Goldston, in *Basic Physical Processes of Toroidal Fusion Plasmas*, Varenna Proceedings, (Monotypia Franchi, Città di Castello, 1985), Vol. 1, p. 165.

¹Work supported by DOE SciDAC GPSC and U.S. DOE Contract No. DE-AC02-76-3073

Gregory Rewoldt
Princeton Plasma Physics Lab.

Date submitted: 20 Jul 2006

Electronic form version 1.4