

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Effects of shear on the magnetic footprint and stochastic layer in double-null divertor tokamak HAMIDULLAH FARHAT, ALKESH PUNJABI, HALIMA ALI, Center for Fusion Research & Training, Hampton University, Hampton, VA 23668 — We have developed a new area-preserving map, called the Adjustable Shear Map, to calculate effects of shear on the magnetic footprint and stochastic layer in double-null divertor tokamak. The map is given by equations $x_{n+1} = x_n - ky_n[(1 - y_n^2)(1 + sy_n) + sx_{n+1}^2]$, $y_{n+1} = y_n + kx_{n+1}[1 + s(x_{n+1}^2 + y_n^2)]$. k is the map parameter and s is the shear parameter. O-point of the map is $(0, 0)$, and the X-points are $(0, 1)$, and $(0, -1)$. For $s=0$, $k=0.6$, the last good surface is $y=0.9918$ with $q \sim 3$. Here we will report on the effects of shear on the stochastic layer and magnetic footprint as the shear parameter is varied from 0 to -1. Here we will report the preliminary results on the effect of shear on the magnetic footprint and the stochastic layer where the shear parameter s has values between -1 and 0. using method of maps [1-4]. This work is done under the DOE grant number DE-FG02-01ER54624. 1. A. Punjabi, A. Boozer, and A. Verma, *Phys. Rev. Lett.*, **69**, 3322 (1992). 2. H. Ali, A. Punjabi, and A. Boozer, *Phys. Plasmas* **11**, 4527 (2004). 3. A. Punjabi, H. Ali, and A. Boozer, *Phys. Plasmas* **10**, 3992 (2003). 4. A. Punjabi, H. Ali, and A. Boozer, *Phys. Plasmas* **4**, 337 (1997).

Halima Ali

Date submitted: 21 Jul 2006

Electronic form version 1.4