

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Alfvén Eigenmodes in Enhanced D-alpha H-mode in Alcator C-Mod¹ J.A. SNIPES, P.T. BONOLI, V. TANG, MIT Plasma Science and Fusion Center, N.N. GORELENKOV, Princeton Plasma Physics Laboratory — Unstable Alfvén eigenmodes are observed in Enhanced D-alpha (EDA) H-mode in Alcator C-Mod at least up to densities of $\bar{n}_e \sim 2.5 \times 10^{20} \text{m}^{-3}$ with strong H-minority ICRF heating. The mode frequencies are typically in the range 400 - 800 kHz. The largest amplitude modes ($\tilde{B}_\theta/B_\theta \sim 5 \times 10^{-5}$ at the wall) often agree in frequency with the expected Toroidal Alfvén Eigenmode (TAE) frequency ($\omega_{TAE} = v_A/2qR$) for $q=1$, indicating that these are core modes. The calculated effective fast ion tail energy ~ 50 keV corresponds to $v_F/v_A \sim 0.4$. These modes are observed to rotate in the electron diamagnetic drift direction, which is opposite to the rotation direction expected for TAEs with a peaked fast ion distribution. Recent calculations with the AORSA and CQL3D codes [1] indicate that the fast ion distribution in C-Mod may be hollow because of poor wave focussing and preferential heating of trapped particles along the resonance. The opposite sign of the gradient of a hollow profile of the fast ion β would then explain the opposite direction of rotation. NOVA-K modelling will be used to compare the stability of L and H-mode conditions and determine the radial structure of these modes.

[1] E. F. Jaeger, et al, to be published in Plasma Physics and Cont. Fus.

¹Supported by USDoE award DE-FC02-99ER54512.

Joseph Snipes
MIT Plasma Science and Fusion Center

Date submitted: 31 Jul 2006

Electronic form version 1.4