

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

High Beta Observations of the Hot Electron Interchange Instability¹ E.E. ORTIZ, M.E. MAUEL, D.T. GARNIER, A.K. HANSEN, Columbia University, J. KESNER, A. BOXER, J.L. ELLSWORTH, I. KARIM, R. BERGMANN, MIT PSFC — High frequency ($f > 1$ MHz) electrostatic fluctuations have been observed in high-beta plasma created in the Levitated Dipole Experiment (LDX). We have previously identified these fluctuations as the Hot Electron Interchange (HEI) instability.² New observations have been made in the presence of the magnetic levitation fields. We find the HEI mode is characterized by frequency sweeping at the drift-resonance of trapped energetic electrons. The fluctuations often appear with coherent structures that have been detected on fast high-impedance electrostatic probes and edge Mirnov sensors. We observe phase shifts using multiple probes that will enable us to determine the toroidal mode number (m) and a higher sampling rate reveals frequency sweeping as high as 40 MHz. Measurements that characterize these modes now incorporate fast magnetic measurements in an attempt to put together a coherent picture of plasma behavior during these modes, including the consequences of these instabilities on plasma formation and pressure limits.

¹This work is supported by U.S. DOE Grants DE-FG02-98ER54458 and DE-FG02-98ER54459

²E. Ortiz, *et al.*, to appear in *J. Fus. Energy*, (2006).

Eugenio Ortiz
Columbia University

Date submitted: 24 Jul 2006

Electronic form version 1.4