

Abstract Submitted
for the DPP07 Meeting of
The American Physical Society

High- β , improved confinement RFP plasmas at high density

M.D. WYMAN, B.E. CHAPMAN, A.F. ALMAGRI, J.K. ANDERSON, D.J. DEN HARTOG, F. EBRAHIMI, D.A. ENNIS, G. FIKSEL, S. GANGADHARA, J.A. GOETZ, R. O'CONNELL, S.P. OLIVA, S.C. PRAGER, J.A. REUSCH, J.S. SARFF, H.D. STEPHENS, UW-Madison, F. BONOMO, P. FRANZ, Consorzio RFX, D.L. BROWER, B.H. DENG, W.X. DING, T. YATES, UCLA, S.K. COMBS, C.R. FOUST, ORNL, D. CRAIG, Wheaton College — In MST discharges with improved confinement, pellet injection has quadrupled the density, with n_e reaching $4 \times 10^{19} \text{m}^{-3}$. The energy confinement time of these high density plasmas is comparable to that at low density, and β_{tot} now reaches 26%. This beta exceeds the Mercier limit for interchange stability, but as yet no indication of interchange is detected experimentally. Beta is also now high enough for global $m = 1$ tearing modes to be linearly unstable, with the instabilities being driven by the pressure gradient rather than the current gradient. Although a β limit has not yet been reached, both the $m = 1$ fluctuation levels and energy transport do increase modestly at high β , suggesting the possibility that beta may eventually be limited by pressure-driven tearing. Work supported by USDOE.

Max Wyman
UW-Madison

Date submitted: 16 Jul 2007

Electronic form version 1.4