

Abstract Submitted  
for the DPP07 Meeting of  
The American Physical Society

**Low Frequency Response of Plasma to MHD Perturbations**<sup>1</sup> M.S. CHU, General Atomics, Y.Q. LIU, Chalmers University — MHD stability of the plasma depends critically on the frequency and wave length of the perturbation. Future tokamaks are expected to operate in regimes where the external macro-scale perturbations have much lower frequencies than the intrinsic dynamical time scales of the particles [1]. This situation calls for a detailed re-examination of the assumptions on previous models of the response of the plasma to MHD perturbations [2]. The kinetic formulation of MHD response [3] is examined numerically in this work. The energy and momentum flux across the plasma surface is expressed in terms of the MHD perturbations. Implication on the stability and plasma response [4] relevant for the resistive wall mode, with its time scale dramatically reduced by the external resistive wall, is discussed.

[1] B. Hu, *et al.*, Phys. Plasmas **12**, 057301 (2005).

[2] A. Bondeson and M.S. Chu, Phys. Plasmas **3**, 3013 (1996).

[3] T.M. Antonsen, Jr. and Y.C. Lee, Phys. Fluids **25**, 132 (1982).

[4] Y.Q. Liu, *et al.*, Phys. Plasmas **7**, 3681 (2000).

<sup>1</sup>Supported by the US DOE under DE-FG03-95ER54309.

M.S. Chu  
General Atomics

Date submitted: 20 Jul 2007

Electronic form version 1.4