

Abstract Submitted
for the DPP07 Meeting of
The American Physical Society

Measurements of core density and magnetic field fluctuations on the MST¹ TRAVIS YATES, WEIXING DING, TROY CARTER, DAVID BROWER, UCLA, JOHN SARFF, STEWART PRAGER, University of Wisconsin, Madison — Fluctuations play an important role in anomalous particle, momentum and energy transport. Core magnetic and density fluctuations are measured using a high-speed, laser-based, Faraday rotation-interferometry system with a bandwidth of 500 kHz and 8 cm chord spacing. Line-averaged measurements of magnetic and density fluctuations can be inverted using a newly developed inversion method to obtain the local spatial profiles. Spatial structure for modes with $m=1$, $n=6$ up to $n=16$, as well as the $m=0$, $n=1$ mode are identified. Fluctuation profiles for modes of given helicity show noticeable changes during the sawtooth cycle. These measurements can also be exploited to determine the local plasma displacement ($\xi_r = \delta n / \nabla n_0$) and radial velocity fluctuations ($\tilde{v}_r = \partial \xi_r / \partial t$) associated with stochastic magnetic fields. Using these parameters, issues related to anomalous particle and momentum transport are addressed. Detailed modeling of local particle density and magnetic fluctuations will be presented.

¹Work supported by U.S. Department Of Energy.

Travis Yates
UCLA

Date submitted: 19 Jul 2007

Electronic form version 1.4