

Abstract Submitted
for the DPP10 Meeting of
The American Physical Society

Modeling laser wakefield accelerators in a Lorentz boosted frame¹

JEAN-LUC VAY, CAMERON GEDDES, Lawrence Berkeley National Laboratory, ESTELLE CORMIER-MICHEL, Tech-X Corporation, DAVID GROTE, Lawrence Livermore National Laboratory, COMPASS TEAM — Modeling of laser-plasma wakefield accelerators in an optimal frame of reference is shown to produce 4-6 orders of magnitude speed-up in calculations from first principles of stages in the 10 GeV-1 TeV energy range, in agreement with the maximum obtainable speedups that are predicted by theory. Obtaining these speedups requires mitigation of a high-frequency instability that otherwise limit effectiveness using a mix of established and novel numerical techniques, and solutions for handling data input and output in a relativistically boosted frame of reference. Using these techniques, agreement at the percentage level is demonstrated between simulations using different frames of reference, with speedups reaching two orders of magnitude, for a 0.1 GeV class stages. The method is then shown to enable direct, efficient full-scale modeling of deeply depleted laser-plasma stages in the 10 GeV-1 TeV range, verifying the scaling of plasma accelerators to very high energies, providing designs for experiments on new lasers such as BELLA.

¹Work supported by US-DOE Contracts DE-AC02-05CH11231 and DE-AC52-07NA27344, and US-DOE SciDAC program COMPASS. Used resources of NERSC and LBNL cluster Lawrencium, supported by US-DOE Contract DE-AC02-05CH11231.

Jean-Luc Vay
Lawrence Berkeley National Laboratory

Date submitted: 08 Jul 2010

Electronic form version 1.4