

Abstract Submitted  
for the DPP10 Meeting of  
The American Physical Society

**A Reduced Model for Predicting Fast-ion  $D_\alpha$  Light in Tokamaks<sup>1</sup>**

L. STAGNER, W.W. HEIDBRINK, UC Irvine, B.A. GRIERSON, PPPL — The fast-ion  $D_\alpha$  (FIDA) diagnostic measures visible photons emitted when energetic deuterium ions undergo charge-exchange reactions with an injected neutral beam [1]. The Doppler shift of the light contains useful information about the fast-ion population but the signal also depends upon many other quantities. A sophisticated forward-modeling code [2] treats all of the relevant atomic physics but there is a need for an approximate code that predicts expected FIDA spectra. This rapid code will be used to remove the FIDA contribution to main-ion  $D_\alpha$  spectra, for data mining, and to guide the execution of experiments. Comparisons of simplified models with full calculations are described.

[1] W.W. Heidbrink et al., Plasma Phys. Control. Fusion **46** (2004) 1855. [2] W.W. Heidbrink et al., Comm. Comp. Phys. (2010) submitted.

<sup>1</sup>Supported by a National Science Foundation Research Experiences for Undergraduates, and the US Department of Energy under DE-FC02-04ER54698 and DE-AC02-09CH11466.

Bill Heidbrink  
UC-Irvine

Date submitted: 15 Jul 2010

Electronic form version 1.4