

Abstract Submitted  
for the DPP11 Meeting of  
The American Physical Society

**Simulation of Observed EGAM Induced Beam-ion Losses in DIII-**

**D**<sup>1</sup> G.J. KRAMER, G.Y. FU, R. NAZIKIAN, Princeton Plasma Physics Laboratory, M.A. VAN ZEELAND, R.K. FISHER, General Atomics, W.W. HEIDBRINK, L. CHEN, UC-Irvine, D.C. PACE, ORISE — In DIII-D a non-perturbative beam-driven instability, the energetic particle geodesic acoustic mode (EGAM) is often observed when counter-going beams are injected in the current ramp-up phase of the discharge. The EGAM is an  $n = 0$  mode in the plasma core with a frequency about 50% below the ideal GAM and it is an excellent mode to validate codes that are used to predict fast-ion losses in burning plasmas. When the EGAM appears the 2.5 MeV neutron emission drops 10% to 15%. This signal is dominated by beam-ion collisions with the thermal plasma and indicates a decrease in the fast-ion population. Recently, accurate measurements have been made of EGAM induced losses with a fast-ion loss detector (FILD) mounted below the mid-plane in DIII-D. We report on state-of-the-art full-orbit following fast-ion loss simulations for the first 50 ms of beam injection using the 3-D beam birth deposition from TRANSP. A very good qualitative agreement between simulation and experiment was found.

<sup>1</sup>Work supported in part by US DOE under DE-AC02-09CH11466, DE-FC02-04ER54698, SC-G903402 and DE-AC05-06OR23100.

G.J. Kramer  
Princeton Plasma Physics Laboratory

Date submitted: 20 Jul 2011

Electronic form version 1.4