

Abstract Submitted
for the DPP12 Meeting of
The American Physical Society

Evolution of Edge Pedestal Profiles Over the L-H Transition¹

M.S. SAYER, W.M. STACEY, J.P. FLOYD, Georgia Institute of Technology, R.J. GROEBNER, General Atomics — The detailed time evolution of thermal diffusivities, electromagnetic forces, pressure gradients, particle pinch and momentum transport frequencies (which determine the diffusion coefficient) have been analyzed during the L-H transition in a DIII-D discharge. Density, temperature, rotation velocity and electric field profiles at times just before and after the L-H transition are analyzed in terms of these quantities. The analysis is based on the fluid particle balance, energy balance, force balance and heat conduction equations, as in Ref. [1], but with much greater time resolution and with account for thermal ion orbit loss. The variation of diffusive and non-diffusive transport over the L-H transition is determined from the variation in the radial force balance (radial electric field, $V \times B$ force, and pressure gradient) and the variation in the interpreted diffusive transport coefficients.

[1] W.M. Stacey and R.J. Groebner, Phys. Plasmas **17**, 112512 (2010).

¹Work supported by the US Department of Energy under DE-FG01-ER54538 and DE-FC02-04ER54698.

Bill Stacey
Georgia Institute of Technology

Date submitted: 12 Jul 2012

Electronic form version 1.4