

Abstract Submitted
for the DPP13 Meeting of
The American Physical Society

Thermal conductivity measurements of CH and Be by refraction-enhanced x-ray radiography¹ YUAN PING, LLNL, JIM KING, OSU, OTTO LANDEN, JEFF KOCH, RUSSELL WALLACE, LLNL, RICK FREEMAN, OSU, TOM BOEHLY, LLE, GLIBERT COLLINS, LLNL — A novel technique, time-resolved refraction-enhanced x-ray radiography, is developed to study thermal conductivity at an interface. Experiments using OMEGA laser have been carried out for CH/Be targets isochorically heated by x-rays to measure the evolution of the density gradient at the interface due to thermal conduction. The sensitivity of this radiographic technique to discontinuities enabled observation of shock/rarefaction waves propagating away from the interface. The radiographs provide enough constraints on the temperatures, densities and scale lengths in CH and Be, respectively. Preliminary data analysis suggests that the thermal conductivities of CH and Be at near solid density and a few eV temperature are higher than predictions by the commonly used Lee-More model. Detailed analysis and comparison with various models will be presented.

¹This project is supported by DOE/OFES Early Career Program. The work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Security, LLC, Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Yuan Ping
LLNL

Date submitted: 09 Jul 2013

Electronic form version 1.4