

Abstract Submitted
for the DPP13 Meeting of
The American Physical Society

The Dynamic Impact of Fusion Self-Heating on Transport Barrier Formation and Control D.E. NEWMAN, Univ. of Alaska Fairbanks, P.W. TERRY, Univ. of Wisconsin Madison, R. SANCHEZ, Univ. Carlos III de Madrid — Over the last 2 decades, simple dynamical models have been able to capture a remarkable amount of the dynamics of the transport barriers found in many devices, including the often disconnected nature of the electron thermal transport channel sometimes observed in the presence of a standard (“ion channel”) barrier. By including in this rich though simple dynamic transport model an evolution equation for electron fluctuations we have investigated the interaction between the formation of the standard ion channel barrier and the somewhat less common electron channel barrier. Barrier formation in the electron channel is even more sensitive to the alignment of the various gradients making up the sheared radial electric field than the ion barrier is. Because of this sensitivity and coupling of the barrier dynamics, the dynamic evolution of the fusion self-heating profile can have a significant impact on the barrier location and dynamics. To investigate this, self-heating has been added to this model and the impact of the self-heating on the formation and controllability of the various barriers is explored. It has been found that the evolution of the heating profiles can suppress or collapse the electron channel barrier leading to the possibility of using NBI for profile/barrier control.

David Newman
Univ. of Alaska Fairbanks

Date submitted: 11 Jul 2013

Electronic form version 1.4