

Abstract Submitted
for the DPP13 Meeting of
The American Physical Society

**ELM-free, Quiescent H-mode Operation in DIII-D Under
Reactor-Relevant Conditions Using Non-Axisymmetric Magnetic Fields¹**

K.H. BURRELL, A.M. GAROFALO, GA, W.M. SOLOMON, PPPL, M.E. FENSTERMACHER, LLNL — Application of static, non-axisymmetric magnetic fields (NAMFs) to DIII-D plasmas allows sustained quiescent H-mode (QH-mode) operation under reactor-relevant conditions of beta, collisionality and torque from neutral beam injection (NBI). QH-mode is an ideal plasma for next step devices, exhibiting H-mode confinement levels while operating without edge localized modes at constant density and radiated power. Peeling-ballooning mode stability theory suggests, and previous studies confirm, that QH-mode operation requires sufficient radial shear in the toroidal rotation near the plasma edge. In past experiments, this rotation shear was predominantly produced by torque from counter-directed NBI. In recent experiments, counter torque due to neoclassical toroidal viscosity produced by the NAMFs gave rise to the necessary edge rotational shear, even overcoming small amounts of co-NBI torque. Experiments in the 2013 campaign have investigated techniques for creating QH-mode plasmas with zero net NBI torque from Ohmic plasmas, opening a path to QH-mode operation with reactor-relevant torque throughout the shot.

¹Work supported by the US DOE under DE-FC02-04ER54698, DE-AC02-09CH11466, and DE-AC52-07NA27344

K.H. Burrell
General Atomics

Date submitted: 11 Jul 2013

Electronic form version 1.4