

Abstract Submitted
for the DPP13 Meeting of
The American Physical Society

Ion Loss as Intrinsic Momentum Source in Tokamaks¹ J.A. BOEDO, UCSD — Measurements of D^+ parallel velocity at the DIII-D edge are consistent with the kinetic loss of thermal ions as the mechanism for edge momentum generation. Edge velocity profiles exhibit a co- I_p peak velocity of 40-60 km/s in OH, L- and H-mode. The flow layer acts as a robust boundary value not affected by NBI injection. D^+ velocity measurements are compared to a first-principles, collisionless, kinetic model predicting the existence of a loss-cone distribution in velocity space resulting in a co- I_p directed velocity. A fine E_r structure, found by probes, has 10-20 kV/m peaks in the scrape-off layer (SOL) and LCFS and when incorporated in the kinetic model, results in: 1) $\sim 30\%$ - 50% increase in the peak parallel velocity over the zero field case and, 2) broadened rotation profile into the SOL. The model-data agreement shows this mechanism is important, competing with pre-sheath acceleration and Pfirsch-Schluter drives. Computations with XGC0, a full-f particle-in-cell drift-kinetic solver with collisional kinetic ions and electrons, and NEO, a drift kinetic code with multiple species and linearized F-P collisions confirms the relevance of the ion orbit loss the impact of kinetic effects on E_r and the measured C^{6+} and D^+ velocities inside the LCFS.

¹Work supported by the US DOE under DE-FG02-07ER54917, DE-FC02-08ER54977, & DE-FC02-04ER54698.

Jose Boedo
UCSD

Date submitted: 12 Jul 2013

Electronic form version 1.4