

Abstract Submitted
for the DPP17 Meeting of
The American Physical Society

Physics of neutral gas jet interaction with magnetized plasmas¹

ZHANHUI WANG, Southwestern Institute of Physics, XUEQIAO XU, Lawrence Livermore National Laboratory, PATRICK DIAMOND, University of California at San Diego, MIN XU, XURU DUAN, DELIANG YU, YULIN ZHOU, YONGFU SHI, LIN NIE, RUI KE, WULV ZHONG, ZHONGBING SHI, AIPING SUN, JIQUAN LI, LIANGHUA YAO, Southwestern Institute of Physics — It is critical to understand the physics and transport dynamics during the plasma fuelling process. Plasma and neutral interactions involve the transfer of charge, momentum, and energy in ion-neutral and electron-neutral collisions. Thus, a seven field fluid model of neutral gas jet injection (NGJI) is obtained, which couples plasma density, heat, and momentum transport equations together with neutrals density and momentum transport equations of both molecules and atoms. Transport dynamics of plasma and neutrals are simulated for a complete range of discharge times, including steady state before NGJI, transport during NGJI, and relaxation after NGJI. With the *trans-neut* module of BOUT++ code, the simulations of mean profile variations and fueling depths during fueling have been benchmarked well with other codes and also validated with HL-2A experiment results. Both fast component (FC) and slow component (SC) of NGJI are simulated and validated with the HL-2A experimental measurements. The plasma blocking effect on the FC penetration is also simulated and validated well with the experiment.

¹This work is supported by the National Natural Science Foundation of China under Grant No. 11575055.

Zhanhui Wang
Southwestern Institute of Physics

Date submitted: 14 Jul 2017

Electronic form version 1.4