

Abstract Submitted
for the DPP17 Meeting of
The American Physical Society

Hamiltonian Single Wave Models to Investigate the Nonlinear Self-Consistent Interaction of Whistler Waves and Electrons¹ CHRIS CRABTREE, GURUDAS GANGULI, ERIK TEJERO, US NRL — We investigate the nonlinear evolution of a self-consistent Hamiltonian model for the interaction of resonant electrons with whistler waves. We find that in the parallel propagating case there are two classes of solutions. The first class has properties similar to previously derived single wave models and involves a perfectly resonant electron beam. We show that test-particle models in the modulated wave-field indicate that the particles are trapped in a first-order island that tracks the location of the majority of particles. We develop a macro-particle model which explains the small frequency and amplitude oscillations. The second class of solutions involves a slightly off-resonant interaction which leads to an amplitude modulation of the wave that resembles the sub-packet structure observed in both chorus and recent laboratory experiments. In the second class, we show that test-particle models demonstrate that instead of being trapped in the primary resonance, particles get trapped in a second-order island chain. The location of the second-order island chain in phase space tracks the location of the majority of electrons. We develop a two macro-particle model which reproduces the amplitude modulation and sub-packet structure of the full model.

¹This work is supported by the Naval Research Laboratory base program and by the National Aeronautics and Space Administration under Grant No. NNH15AZ90I.

Chris Crabtree
US NRL

Date submitted: 14 Jul 2017

Electronic form version 1.4