

Abstract Submitted  
for the DPP17 Meeting of  
The American Physical Society

**Testing low-mode symmetry control with low-adiabat, extended pulse-lengths in BigFoot implosions on the National Ignition Facility**  
MATTHIAS HOHENBERGER, D. T. CASEY, C. A. THOMAS, K. L. BAKER, B. K. SPEARS, S. F. KHAN, O. A. HURRICANE, D. CALLAHAN, Lawrence Livermore Natl Lab — The Bigfoot approach to indirect-drive inertial confinement fusion (ICF) has been developed as a compromise trading high-convergence and areal densities for high implosion velocities, large adiabats and hydrodynamic stability. Shape control and predictability are maintained by using relatively short laser pulses and merging the shocks within the DT-ice layer. These design choices ultimately limit the theoretically achievable performance, and one strategy to increase the 1-D performance is to reduce the shell adiabat by extending the pulse shape. However, this can result in loss of low-mode symmetry control, as the hohlraum “bubble,” the high-Z material launched by the outer-cone beams during the early part of the laser pulse, has more time to expand and will eventually intercept inner-cone beams preventing them from reaching the hohlraum waist, thus losing equatorial capsule drive. We report on experimental results exploring shape control and predictability with extended pulse shapes in BigFoot implosions. Prepared by LLNL under Contract DE-AC52-07NA27344.

Matthias Hohenberger  
Lawrence Livermore Natl Lab

Date submitted: 16 Jul 2017

Electronic form version 1.4