

Abstract Submitted
for the DPP19 Meeting of
The American Physical Society

Study of argon impurity transport by X-ray imaging crystal spectrometer on J-TEXT WEI YAN, ZHONGYONG CHEN, Huazhong University of Science and Technology, XIAOLONG ZHANG, Shenzhen University, ZHIFENG CHENG, Huazhong University of Science and Technology, JIANCHAO LI, Shenzhen University, J-TEXT TEAM — A tangential X-ray imaging crystal spectrometer (XICS) has been upgraded on J-TEXT tokamak to measure the electron/ion temperature and the plasma toroidal rotation velocity. The XICS has been designed to receive emissions of Ar XVII from -13 cm to $+13$ cm region with a spatial resolution of 1.8 cm in the vertical direction. The temporal evolution of argon impurity density profiles after an argon gas puff could be observed with a time resolution of up to 2 ms. The emissions of Ar XVII can be modulated by the resonant magnetic perturbations (RMPs) which indicates that the transport of argon is affected by the RMPs significantly. The 2/1 RMPs can lead to field penetration with enough RMPs amplitude. The XICS provides a tool for the study of the transport of argon impurities during the penetration of RMPs. During the field penetration phase, the emissions of Ar XVII decreased. The phenomena show that the transport of argon impurity at $r/a \sim 0.5$ has been enhanced during the field penetration phase. The Ar XVII profile can be changed by the different 2/1 RMP phases. STRAHL can give the ratio between D and v profiles for the argon. The result shows that the argon transport can be affected by 2/1 RMP and its phase.

Wei Yan
Huazhong University of Science and Technology

Date submitted: 10 Jul 2019

Electronic form version 1.4