

Abstract Submitted  
for the HAW14 Meeting of  
The American Physical Society

**Transverse dynamics of multi-strange hadrons** SHIORI TAKEUCHI, Sophia University, KOICHI MURASE, The University of Tokyo, TETSUFUMI HIRANO, Sophia University, PASI HOUVINEN, Johann Wolfgang Goethe-Universität, YASUSHI NARA, Akita International University — We study the hadronic rescattering effects, in particular for multi-strange hadrons, on the final observables within an integrated dynamical approach in which a (3+1)D ideal hydrodynamic model is combined with hadronic cascade model, JAM. Since the measured observables contain all the information throughout the space-time evolution in relativistic heavy ion collisions, dissipation caused by hadronic rescattering in the late stage must be taken into account to investigate the properties of the quark gluon plasma. By comparing final observables to those obtained just after the QGP stage in this approach, we quantify the hadronic rescattering effects on particle ratio, azimuthal anisotropic flow  $v_n$  and mean transverse momentum  $\langle p_T \rangle$ . Furthermore we show that multi-strange hadrons can be direct probes of the QGP at hadronization due to a fact that multi-strange hadrons rescatter less than non-strange hadrons.

Shiori Takeuchi  
Sophia University

Date submitted: 16 Jun 2014

Electronic form version 1.4