

Abstract Submitted
for the MAR05 Meeting of
The American Physical Society

Korringa-Like Nuclear Spin-Lattice Relaxation in a 2DES at $\nu = 1/2$ L.A. TRACY, I.B. SPIELMAN, J.P. EISENSTEIN, Caltech, L.N. PFEIFFER, K.W. WEST, Bell Labs, Lucent Technologies — Via a resistively-detected NMR technique, the nuclear spin lattice relaxation time T_1 of ^{71}Ga at low temperatures has been measured in a GaAs/AlGaAs heterostructure containing two weakly-coupled 2D electron systems (2DES), each at Landau level filling $\nu = 1/2$. Incomplete electronic spin polarization, which has been reported previously [1,2] for low density 2DESs at $\nu = 1/2$, should facilitate hyperfine-coupled nuclear spin relaxation owing to the presence of both electron spin states at the Fermi level. Within composite fermion theory, a Korringa law temperature dependence: $T_1 T = \text{constant}$, is expected for temperatures $T < 1$ K. Our measurements made at temperatures in the range $35 \text{ mK} < T < 200 \text{ mK}$, show that T_1 rises less rapidly with falling temperature than this law predicts. This may suggest the existence of alternate nuclear spin relaxation mechanisms in this system.

[1] I. V. Kukushkin, K. v. Klitzing, and K. Eberl. Phys. Rev. Lett. 82, 3665 (1999); A. E. Dementyev, et al., Phys. Rev. Lett. 83, 5074 (1999); S. Melinte, et al., Phys. Rev. Lett. 84, 354 (2000).

[2] I.B. Spielman, L.A. Tracy, J.P. Eisenstein, L.N. Pfeiffer, K.W. West, condmat/0410092.

This work was supported by the DOE, NSF, and NDSEG.

Lisa Tracy
Caltech

Date submitted: 29 Nov 2004

Electronic form version 1.4