

Abstract Submitted
for the MAR05 Meeting of
The American Physical Society

Potential-Energy Surface of Infinite Helical Polypeptides

JOEL IRETA, MATTHIAS SCHEFFLER, Fritz-Haber-Institut der Max-Planck-Gesellschaft — The potential-energy surfaces of infinite polyalanine and polyglycine chains in helical conformation are studied using density-functional theory in the Perdew, Burke and Ernzerhof approximation to the exchange-correlation functional (DFT-PBE). Minima associated to a π -helix, α -helix and 3_{10} -helix conformations are identified for both polypeptides. For polyalanine the α -helix minimum is the lowest in energy. However for polyglycine π -helix and α -helix minima are degenerated within the DFT accuracy. The α -helix is found to undergo structural transitions to a π - or 3_{10} -helix when the length of the helix is strained by more than 10%. The barriers for the structural transitions mainly associated to the breaking of the hydrogen bonds are considerably affected by the side group in polyalanine. We find this effect can not be solely attributed to repulsive interactions between the side group and the helix backbone but to sizeable changes in covalent bonds in the peptide unit of polyalanine with respect to polyglycine.

Joel Ireta
Fritz-Haber-Institut der Max-Planck-Gesellschaft

Date submitted: 30 Nov 2004

Electronic form version 1.4