

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Atomistic Studies of Plastic Deformation and Dissipation in Crystalline HMX EUGENIO JARAMILLO, THOMAS D. SEWELL, Los Alamos National Laboratory, ALEJANDRO STRACHAN, Purdue University — We are using large scale molecular dynamics simulations of crystalline octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX) to better understand the dominant fundamental mechanisms of inelastic deformation and other dissipative processes in anisotropic organic molecular crystals. A fully flexible force field (Smith, G. D. and Bharadwaj, R. K.; *J. Phys. Chem. B* 1999, 103, 3570) used in numerous preceding studies is used without modification in the present work. Our results, based on the results of simulations containing 25,000-250,000 molecules, indicate a large degree of directional anisotropy in response to compression, for both quasi-static and shock loading. Plastic deformation is observed for some loading directions whereas solid-solid phase transitions are observed for others. The emphasis of the present talk will be identifying and characterizing detailed molecular mechanisms and rate dependencies in those cases for which dislocation-induced plasticity occurs.

Eugenio Jaramillo
Los Alamos National Laboratory

Date submitted: 27 Oct 2005

Electronic form version 1.4