

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Magnetic phase separation in electron-doped Bi_{1-x}Ca_xMnO₃ systems YUHAI QIN, TREVOR A. TYSON, New Jersey Institute of Technology, KENJI SHIMIZU, Toyama University — The manganite system Bi_{1-x}Ca_xMnO₃ possesses intriguing properties in the low bismuth doping region. In this electron doped region ($0.6 < x < 1$), a ferromagnetic (FM) moment of ~ 1.2 Bohr magnetons per Mn site is found for $x \sim 0.875$. The magnetic moment per Mn site maintains a value $\sim 1/3$ the theoretical limit even in fields as high as 60 T. The physical origin of this high moment region is not well understood. Various models including canted ferromagnetism and ferromagnetic clusters hosted by an antiferromagnetic background have been proposed. In our previous work, we have conducted small-angle neutron scattering (SANS) on Bi_{0.125}Ca_{0.875}MnO₃ polycrystalline samples as has revealed existence of FM clusters embedded in an AFM background. New ⁵⁵Mn NMR results give more evidence supporting of this heterogeneous phase model: resonance signals from both AFM and FM phases were identified. More progress from multiple-temperature Bi-L₃ edge XAFS measurements will be presented as well. This work is supported by NSF DMR-0209243 and NSF DMR-0512196.

Yuhai Qin
New Jersey Institute of Technology

Date submitted: 23 Nov 2005

Electronic form version 1.4