

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

**Conductance Fano lineshapes for Kondo impurities on surfaces:
A numerical renormalization group description.** NANCY SANDLER, LUIS
DIAS DA SILVA, SERGIO ULLOA, Department of Physics and Astronomy, Ohio
University — Scanning tunneling microscopy (STM) measurements of Kondo im-
purities on metallic surfaces has been an active field in recent years. For a flat
density-of-states (DoS) near the Fermi energy in the host metal, the low-bias STM
conductance acquires the characteristic Fano lineshape, with width proportional to
the Kondo temperature T_K . In this work, we study how this picture is modified
when a *structured* DoS (non-flat) is considered. A variety of physical effects can
introduce peak/dips in the DoS, including the presence of a second impurity, hy-
bridization between surface and bulk conduction states, and a magnetic impurity
embedded in a molecule. Using numerical renormalization group techniques, we cal-
culate the low-temperature conductance for this system. The zero-bias dip in the
Fano conductance is modified by the presence of resonances or anti-resonances in
the DoS near E_F . In particular, for DoS with pseudogaps and impurities in the
mixed-valence regime, zero-bias Fano-like dips appear *even when no Kondo state
has developed*, but governed by energy scales much larger than T_K . We further show
that measurements of the scattering phase could be used as an additional probe into
the Kondo regime. Supported by NFS-NIRT.

Nancy Sandler
Department of Physics and Astronomy, Ohio University

Date submitted: 27 Nov 2005

Electronic form version 1.4