

Abstract Submitted  
for the MAR06 Meeting of  
The American Physical Society

**Fermi Surface of the Half Heusler Compounds  $\text{Ce}_{1-x}\text{La}_x\text{BiPt}$  A.**

D. BIANCHI, J. WOSNITZA, HLD, FZR, P.O. Box 51 01 19, D-01314 Dresden, Germany, N. KOZLOVA, D. ECKERT, L. SCHULTZ, I. OPHALE, S. ELGAZZAR, M. RICHTER, IFW Dresden, D-01171 Dresden, Germany, J. HAGEL, M. DOERR, IFP, TU Dresden, D-01062 Dresden, Germany, G. GOLL, H. V. LÖHNEYSEN, Physikalisches Institut, Universität Karlsruhe, D-76128 Karlsruhe, Germany, G. ZWICKNAGL, Institut für Mathematische Physik, Technische Universität Braunschweig, D-38106 Braunschweig, Germany, T. YOSHINO, T. TAKABATAKE, Department of Quantum Matter, ADSM, Hiroshima University, Higashi-Hiroshima 739-8530, Japan — We report on the Fermi surface in the correlated half-Heusler compound  $\text{Ce}_{1-x}\text{La}_x\text{BiPt}$ . In  $\text{CeBiPt}$  we find a field-induced change of the electronic band structure as discovered by electrical-transport measurements in pulsed magnetic fields. For magnetic fields above  $\sim 25$  T, the charge-carrier concentration determined from Hall-effect measurements increases nearly 30%, whereas the Shubnikov–de Haas (SdH) signal disappears at the same field. In the non- $4f$  compound  $\text{LaBiPt}$  the Fermi surface remains unaffected, suggesting that these features are intimately related to the Ce  $4f$  electrons. Electronic band-structure calculations point to a  $4f$ -polarization-induced change of the Fermi-surface topology. In order to test this hypothesis, we have measured the (SdH) signal in a  $\text{Ce}_{0.95}\text{La}_{0.05}\text{BiPt}$  sample with a low La concentration.

A. D. Bianchi  
FZR/HLD

Date submitted: 30 Nov 2005

Electronic form version 1.4