

Abstract Submitted  
for the MAR07 Meeting of  
The American Physical Society

**Nonadiabatic MD simulations of  $\text{IBr}^-(\text{CO}_2)_n$  photodissociation**  
MATTHEW A. THOMPSON, JOSHUA MARTIN, JOSHUA DARR, JACK BARBERA, VLADIMIR DRIBINSKI, W. CARL LINEBERGER, ROBERT PARSON, JILA and Department of Chemistry, University of Colorado at Boulder — Potential energy curves for the ground and valence excited states of  $\text{IBr}^-$  have been calculated at the MRCI level using the MOLPRO *ab initio* package. Spin-orbit coupling was calculated via a spin-orbit ECP. Charge densities, transition moments, and nonadiabatic coupling matrix elements constructed from a distributed multipole analysis of the *ab initio* wavefunctions<sup>1</sup> were then used to carry out nonadiabatic molecular dynamics simulations of the photodissociation of  $\text{IBr}^-$  in  $\text{CO}_2$  clusters. Experimental studies have demonstrated a large variation in ground-state recombination times which are supported by our simulations. We propose a mechanism of excited-state trapping and a configurational transition state which leads to recombination times on the order of 10-20 ps for  $n=5,13$  up to 1 ns for  $n=8,10$ .

<sup>1</sup>Maslen, Faeder and Parson, Molecular Physics, 1998

Matthew Thompson  
JILA and Department of Chemistry, University of Colorado at Boulder

Date submitted: 10 Nov 2006

Electronic form version 1.4