

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

First-Principles Studies of sila-Diamondoids¹ STEVEN L. RICHARDSON, Howard University, RAJENDRA R. ZOPE, TUNNA BARUAH, University of Texas at El Paso, MARK R. PEDERSON, Naval Research Laboratory — While there has been recent progress in the isolation of carbon-based diamondoids from petroleum oil by the Chevron-Texaco group, there is still no experimental evidence for the existence of the silicon analogue of adamantane, sila-adamantane ($\text{Si}_{10}\text{H}_{16}$). As adamantane forms the central building block for diamondoids, we speculate that sila-adamantane could serve as the template for a novel class of materials known as sila-diamondoids. We predict that because sila-diamondoids are nanostructures derived from bulk crystalline silicon they will have important applications in molecular electronics and nanotechnology. Recently, Fischer, Baumgartner, and Marschner (*Science* **310**, 825 (2005)) have made an important contribution to the field of sila-diamondoids with the synthesis of a four-fold silylated molecule, $\text{C}_{24}\text{H}_{72}\text{Si}_{14}$. In this work, we show that density-functional theory is capable of calculating the structural, electronic, and vibrational properties of $\text{C}_{24}\text{H}_{72}\text{Si}_{14}$ that compare very well with the recent experimental data.

¹This work is supported by NSF CREST Grant No. HRD-0317607.

Steven L. Richardson
Howard University

Date submitted: 21 Nov 2006

Electronic form version 1.4