

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Frustration in the Hubbard model: a quantum cluster study ANDRIY NEVIDOMSKYY, University of Sherbrooke, Quebec, Canada, CHRISTIAN SCHEIBER, Technical University of Graz, Austria, DAVID SENECHAL, University of Sherbrooke, Quebec, Canada — The role of frustration in the Hubbard model is studied on the square lattice with nearest and next-nearest neighbour hoppings t and t' using the Variational cluster perturbation theory (VCPT, see [1]). We find two phases with long-range magnetic order: the usual antiferromagnet (AF1) phase, stable at small t'/t , and the so-called superantiferromagnetic phase (AF2) for large frustration. These are separated by a phase with no magnetic order. We also find d-wave superconductivity ($d_{x^2-y^2}$) for small values of $U \leq 4t$ and sufficiently weak frustration. The Mott-Hubbard transition is discussed in this context. We also compare the classical phase diagram obtained from the large- U expansion with that of the frustrated J_1 - J_2 Heisenberg model.

[1] M. Potthoff, M. Aichhorn, C. Dahnken, Phys. Rev. Lett. **91**, 206402 (2003).

Andriy Nevidomskyy
University of Sherbrooke, Quebec, Canada

Date submitted: 22 Nov 2006

Electronic form version 1.4