

Abstract Submitted  
for the MAR07 Meeting of  
The American Physical Society

**A non-Griffiths-like clustered phase above the Curie temperature of the doped perovskite cobaltite  $\text{La}_{1-x}\text{Sr}_x\text{CoO}_3$** <sup>1</sup> CHUNYONG HE, MARIA TORIJA, JING WU, University of Minnesota, JEFF LYNN, NIST Center for Neutron Research, JOHN MITCHELL, Argonne National Laboratory, CHRIS LEIGHTON, University of Minnesota — The existence of preformed clusters above the  $T_c$  of the doped perovskite manganites is well established and, in many cases, conforms to the expectations for a Griffiths phase. We show here that the phase-separated perovskite cobaltite ( $\text{La}_{1-x}\text{Sr}_x\text{CoO}_3$ ) also exhibits a clustered state above the  $T_c$  in the ferromagnetic phase. The formation of magnetic clusters at a well-defined temperature ( $T^*$ ) is revealed in the small-angle neutron scattering, d.c. susceptibility, and resistivity. Remarkably, this clustered state has none of the characteristics of a Griffiths phase; the deviation from Curie-Weiss behavior is opposite to expectations and is not field dependent, and  $T^*$  does not correspond to the undiluted  $T_c$ . These results demonstrate that although the Griffiths phase may occur in many systems with quenched disorder, it is not universally applicable to the randomly doped transition metal oxides.

<sup>1</sup>We acknowledge support from DOE (DE-FG02-06ER46275), NSF (DMR-0509666), and the ACS PRF, as well as neutron facilities at NIST. Work at ANL supported by US DoE under W-31-109-ENG-38.

Chunyong He  
University of Minnesota

Date submitted: 20 Nov 2006

Electronic form version 1.4