

Abstract Submitted  
for the MAR07 Meeting of  
The American Physical Society

**Angular dependent magnetorestriction study of CeCoIn<sub>5</sub>**  
CATALIN MARTIN, VICTOR. F. CORREA, KENNETH M. PURCELL, TIMOTHY P. MURPHY, ERIC C. PALM, STANLEY W. TOZER, National High Magnetic Field Laboratory, Florida State University, Tallahassee, Florida 32310, USA, GEORGE M. SCHMIEDESHOFF, Department of Physics, Occidental College, Los Angeles, California, 90041, J. C. COOLEY, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA, S. W. TOZER TEAM, G. M. SCHMIEDESHOFF COLLABORATION, J. C. COOLEY COLLABORATION — Using a capacitive dilatometer designed to operate on the rotating stage of a top loading dilution refrigerator probe, we measured the magnetorestriction  $(L(H) - L(0))/L(0)$ , of the Heavy Fermion Superconductor CeCoIn<sub>5</sub>, along all three crystallographic direction, and with the magnetic field applied parallel both to the superconducting *ab*-planes and to the *c*-axis. When  $B||ab$ , the *c*-axis magnetorestriction features, besides a sharp jump at the upper critical field, a continuous, second-order like, transition at a lower field, which is consistent with previous observations from specific heat measurements. Our results are consistent with a field induced modulation of the superconducting order parameter (FFLO state), rather than with a field induced magnetic ordering.

Catalin Martin  
National High Magnetic Field Laboratory, Florida State University,  
Tallahassee, Florida 32310, USA

Date submitted: 01 Dec 2006

Electronic form version 1.4