

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

**Fluctuations in Two-Dimensional
Superconducting NbN Nanobridges and Nanostructures Meanders**

JENNIFER KITAYGORSKY, I. KOMISSAROV, A. JUKNA, University of Rochester, O. MINAEVA, N. KAUROVA, A. DIVOCHIY, A. KORNEEV, M. TARKHOV, B. VORONOV, I. MILOSTNAYA, G. GOL'TSMAN, Moscow State Pedagogical University, R. SOBOLEWSKI — We have observed fluctuations, manifested as sub-nanosecond to nanosecond transient, millivolt-amplitude voltage pulses, generated in two-dimensional NbN nanobridges, as well as in extended superconducting meander nanostructures, designed for single photon counting. Both nanobridges and nano-stripe meanders were biased at currents close to the critical current and measured in a range of temperatures from 1.5 to 8 K. During the tests, the devices were blocked from all incoming radiation by a metallic enclosure and shielded from any external magnetic fields. We attribute the observed spontaneous voltage pulses to the Kosterlitz-Thouless-type fluctuations, where the high enough applied bias current reduces the binding energy of vortex-antivortex pairs and, subsequently, thermal fluctuations break them apart causing the order parameter to momentarily reduce to zero, which in turn causes a transient voltage pulse. The duration of the voltage pulses depended on the device geometry (with the high-kinetic inductance meander structures having longer, nanosecond, pulses) while their rate was directly related to the biasing current as well as temperature.

Jennifer Kitaygorsky
University of Rochester

Date submitted: 20 Nov 2006

Electronic form version 1.4