

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Low temperature specific heat of geometrically frustrated $\text{Gd}_2\text{Sn}_2\text{O}_7$ ¹ KATE ROSS, JEFFREY QUILLIAM, CHAS MUGFORD, University of Waterloo, LINTON CORRUCINI, University of California, Davis, JAN KYCIA, University of Waterloo — Previous measurements of the specific heat of $\text{Gd}_2\text{Sn}_2\text{O}_7$ showed a T^2 power law below a strongly first-order phase transition, though there was some indication of a deviation from this power law below 500 mK². Theory has predicted that anisotropy due to the dipolar interaction leads to a gapped spin wave spectrum resulting in an exponential specific heat as T approaches 0.³ We will present specific heat measurements to below 100 mK. Preliminary results show a deviation from the T^2 law which supports the above theoretical model.

¹Work was supported by NSERC, CFI, MMO, EMK, The Research Corporation

²P. Bonville *et al.* *J. Phys.: Cond. Mat.* **15**, 7777 (2003).

³A. G. Del Maestro and M. J. P. Gingras, *J. Phys.: Cond. Mat.* **16**, 3339 (2004).

Kate Ross
University of Waterloo

Date submitted: 20 Nov 2006

Electronic form version 1.4