

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

**Total energy calculations using the transcorrelated method:
wavefunction approach for solids** KEITARO SODEYAMA, REI SAKUMA,
SHINJI TSUNEYUKI, Department of Physics, University of Tokyo — To calcu-
late the electronic structures of solids including electron correlation effects, we have
developed the transcorrelated (TC) method which was first proposed by Boys and
Handy. In this method, the wave function is represented by a correlated wave
function $F\Phi$, where Φ is a single Slater determinant and F is a Jastrow func-
tion, $F = \exp[-\sum_{i<j} u_{ij}]$. u_{ij} is a two-body function called Jastrow factor. The
many-body Hamiltonian H is similarity transformed to an effective Hamiltonian
 $H_{TC} = F^{-1}HF$ with up-to-three-body interaction. One-electron orbitals and or-
bital energies are optimized by solving a set of Hartree-Fock(HF)-like equations
derived by minimizing the variance of the H_{TC} . In this study, the total energies of
crystalline silicon and carbon are calculated by the TC methods at various lattice
constants. The bulk moduli of silicon by HF and TC methods were 115 and 102 GPa.
Those of carbon were 524 and 501 GPa. Comparing with the experimental values
of 99.2 and 443 GPa, the TC method, which includes the electron correlation effect
in the framework of wavefunction approach, improves the HF results.

Keitaro Sodeyama
Department of Physics, University of Tokyo

Date submitted: 28 Dec 2006

Electronic form version 1.4