

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Absence of Hole Confinement in Transition Metal Oxides with Orbital Degeneracy¹ MARIA DAGHOFER, University of Tennessee, KRZYSZTOF WOHLFELD, ANDRZEJ M. OLES, Jagellonian University, ENRICO ARRIGONI, Graz University of Technology, PETER HORSCH, Max-Planck-Institut FKF — The compounds with orbital degrees of freedom exhibit many possible scenarios for hole propagation which in most cases lead to hole localization [1]. Here we investigate the spectral properties of a hole moving in a two-dimensional Hubbard model for strongly correlated t_{2g} electrons. Although superexchange interactions are Ising-like, a quasi-one-dimensional coherent hole motion arises due to effective three-site terms. This mechanism is fundamentally different either from the hole motion via quantum fluctuations in the conventional spin model with SU(2) symmetry or from the e_g orbital model [2]. The present orbital model describes also propagation of a hole in some e_g compounds [3], and we argue that orbital degeneracy alone does not lead to hole self-localization. [1] J. Zaanen and A.M. Oleś, Phys. Rev. B **48**, 7197 (1993). [2] J. van den Brink, P. Horsch, and A.M. Oleś, Phys. Rev. Lett. **85**, 5174 (2000). [3] M. Daghofer, A.M. Oleś, and W. von der Linden, Phys. Rev. B **70**, 184430 (2004).

¹We acknowledge support by Foundation for Polish Science (FNP), by the Polish Ministry of Science and Education Project No. N202 068 32/1481, and by the Austrian Science Fund (FWF Project P18505-N16).

Andrzej M. Oles
Jagellonian University

Date submitted: 02 Dec 2007

Electronic form version 1.4