

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

**Viscosity contribution to the impurity resistivity of metals by means of the current-density functional theory** VLADIMIR U. NAZAROV, Research Center for Applied Sciences, Academia Sinica, Taiwan, GIOVANNI VIGNALE, University of Missouri, Columbia, USA — Within the time-dependent density functional theory formalism we relate the impurity resistivity  $\rho$  of a metal to the friction coefficient  $Q$  of the metal for the same impurity moving with the infinitesimally small velocity, i.e.,  $\rho = n_i Q / n_e^2$  (1), where  $n_i$  and  $n_e$  are the concentrations of the randomly distributed impurities and the valence electrons, respectively. While Eq.(1) occurs trivial within the single-particle theory with the scattering at the *statically* screened impurities, its general validity within the many-body theory with the *dynamical* exchange and correlation included presents a progress. We utilize results [1,2] on  $Q$  of the electron liquid to put the electron-electron scattering contribution into the terms of the viscosity coefficients [3]. Calculations of the residual resistivity of aluminum as a function of the atomic number of the impurity are performed, improving the agreement with experiment compared to the single-particle theory [4]. [1].V. U. Nazarov, J. M. Pitarke, C. S. Kim, and Y. Takada, Phys. Rev. B **71**, 121106(R) (2005). [2].V. U. Nazarov, J. M. Pitarke, Y. Takada, G. Vignale, and Y.-C. Chang, Phys. Rev. B **76**, 205103 (2007). [3].G. Vignale, C. A. Ullrich, and S. Conti, Phys. Rev. Lett. **79**, 4878 (1997). [4].M. J. Puska and R. M. Nieminen, Phys. Rev. B **27**, 6121 (1983).

Vladimir U. Nazarov  
Research Center for Applied Sciences, Academia Sinica, Taiwan

Date submitted: 21 Nov 2007

Electronic form version 1.4