

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

**Cavity cooling of  $^{88}\text{Sr}^+$**  DAVID LEIBRANDT, YAT SHAN AU, ISAAC CHUANG, MIT — Cavity cooling is a method of laser cooling which uses coherent scattering to cool atoms [V. Vuletić and S. Chu, PRL **84**, 3787 (2000)]. The closed atomic transition used in Doppler cooling is replaced by a cavity resonance, so cavity cooling can be used to cool to sub-Doppler temperatures and is in principle applicable to complicated atoms or molecules without closed transitions. We describe an experiment to study three-dimensional cavity cooling of a single  $^{88}\text{Sr}^+$  ion confined in a linear RF Paul trap. Large cooling rates can be attained by operating near the 422 nm  $S_{1/2} \leftrightarrow P_{1/2}$  optical dipole transition and using a 5 cm long near-confocal Fabry-Pérot cavity with commercially available mirrors of finesse  $10^4$ . Given a cavity alignment error  $\leq 10 \mu\text{m}$  and a trap frequency of 1 MHz, the resolved sideband cavity cooling limit is  $\leq 5$  motional quanta. We present details of the experimental proposal and its implementation.

David Leibbrandt  
MIT

Date submitted: 24 Nov 2007

Electronic form version 1.4