

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Optical properties of ErH_{2+x} : First principles calculations and experimental measurements.¹ CLARK SNOW, THOMAS MATSSON, Sandia National Laboratories — Rare earth and transition metal hydrides exhibit many interesting physical phenomena, from metal-semiconductor transitions to transparency changes at RT as a function of hydrogen content. Electrical resistivity measurements by P. Vajda (1) indicate that ErH_{2+x} undergoes a metal-semiconductor transition between 240-290K and an antiferromagnetic transition between 1.75-2.3K depending on hydrogen content. These same transitions should also cause profound changes in the optical properties. This work will present first principles calculations of the optical properties of ErH_{2+x} from 300-600K where x ranges from -0.2 to 0.2. The calculations will be compared to experimental results on thin films of ErH_{2+x} which were obtained as a function of temperature from 10-600K and hydrogen content.

[1] P. Vajda and J.N. Daou, Phys. Rev. B., Vol. 49, p. 3275 (1994).

¹Sandia is a multi-program laboratory operated by Sandia Corporation, a Lockheed Martin Company, for the United States Department of Energy's National Nuclear Security Administration under Contract DE-AC04-94AL85000.

Clark Snow
Sandia National Laboratories

Date submitted: 12 Dec 2007

Electronic form version 1.4