

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

The condensation phenomena of conserve-mass aggregation model with mass-dependent fragmentation DONG-JIN LEE, SUNGCHUL KWON, YUP KIM, Department of Physics and Research Institute for Basic Sciences, Kyung Hee University — We study a conserved mass aggregation model with mass-dependent fragmentation in regular lattice and scale-free networks. In the model, the whole mass m of a site isotropically diffuse with unit rate. With rate ω , a mass m^λ is fragmented from the site and moves to a randomly selected nearest neighbor site. Since the fragmented mass is smaller than the whole mass m of a site for $\lambda < 1$, the on-site attractive interaction exists for the case. For $\lambda = 0$, the model is known to undergo the condensation phase transitions as the density of total masses (ρ) increases beyond a critical density ρ_c . For $0 < \lambda < 1$, we numerically confirm for several values of ω that ρ_c diverges with the system size L . Hence in thermodynamic limit, the condensed phase disappears and no transitions take place in one dimension. We also explain that there are no transitions in any dimension. On scale-free networks with degree distribution $P(k) \sim k^{-\gamma}$, we numerically confirm for $\gamma > 3$ that the condensation transitions occurs at $\rho_c > 0$ and its nature is the same as that in regular lattice. However, for $\gamma \leq 3$, the condensation always takes place for $\lambda < \lambda_c$ and masses distribute uniformly without aggregation for $\lambda \geq \lambda_c$. We derive $\lambda_c = 1/\gamma - 1$ via mean-field argument.

Dong-Jin Lee
Department of Physics and Research Institute for
Basic Sciences, Kyung Hee University

Date submitted: 01 Dec 2007

Electronic form version 1.4