

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

**A Mathematical Model for Calculating the Effect of Toroidal Geometry on the Measured Magnetic Field**<sup>1</sup> BRENDA SKOCZELAS, RANJITH WIJESINGHE, Ball State University — A mathematical model to calculate the measured magnetic field from a stimulated nerve has been presented in the past. Traditionally, electrodes have been used to measure these propagating action signals in nerves, but a less invasive technique is to use toroids. However, up until now, when using a toroidal transformer to record the nerve action currents, the thickness of the toroid has yet to be considered in the model and how it may affect the propagating compound action potential. In this presentation, we will discuss the development of a new model, to which the thickness of the toroid is taken into account. These dimensions are important because the toroid represents an inhomogeneity in the extracellular medium that redistributes the extracellular current. In the past, toroids with very small diameters have been used and as they may not disrupt the action current. With a better understanding of the toroidal effects, we may be able to increase the accuracy and dependency of such measured magnetic signals. The final goal will be to compare our theoretical model to experimentally gathered data.

<sup>1</sup>Special thanks to Prof. John P. Wikswo of Vanderbilt University.

Brenda Skoczelas  
Ball State University

Date submitted: 03 Dec 2007

Electronic form version 1.4