

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Magnetic excitations in a quantum spin dimer system $\text{Ba}_3\text{Cr}_2\text{O}_8$
MAIKO KOFU, JUNG-HWA KIM, SEUNG-HUN LEE, Univ. of Virginia, BELLA LAKE, Hahn-Meitner Institute, YIMING QIU, National Institute of Standards and Technology, HIROAKI UEDA, YUTAKA UEDA, ISSP, Univ. of Tokyo — We report our neutron scattering measurements on a powder sample of a new quantum spin system $\text{Ba}_3\text{Cr}_2\text{O}_8$ in which Cr^{5+} ($3d^1$, $S = 1/2$) ions form a double-layered triangular lattice. The system does not undergo any magnetic ordering down to 50 mK. Instead, bulk susceptibility data exhibit a broad peak around 16K. Our inelastic neutron scattering data at temperatures lower than 30 K show a prominent excitation at $\hbar\omega = 2.2$ meV. Q- and temperature dependences of the integrated intensity of the excitation can be well accounted for by a model of weakly coupled quantum dimers. When an external magnetic field is applied, the excitation splits into three energies, which confirms the dimer model. We have investigated how the singlet-to-triplet excitations evolve with increasing the magnetic field up to 14.5 Tesla.

Maiko Kofu
Univ. of Virginia

Date submitted: 27 Nov 2007

Electronic form version 1.4