

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Interorbital Pairing and its Physical Consequences in Iron Pnictide Superconductors¹ YI GAO, WU-PEI SU, Department of Physics and Texas Center for Superconductivity, University of Houston, JIAN-XIN ZHU, Theoretical Division, Los Alamos National Laboratory — We study interorbital pairings of iron pnictide superconductors within a minimal two-orbital tight-binding model. We find that in real space, a set of self-consistently determined pairing order parameters forms two sublattices with a relative phase of π and the pairing symmetry is $d_{x^2-y^2} \sim \cos k_x - \cos k_y$. In momentum space, it corresponds to the η pairing proposed by C. N. Yang [Phys. Rev. Lett. **63**, 2144 (1989)], with nonzero momenta of Cooper pairs. One physical consequence of this type of pairing is the existence of a significant amount of zero energy (gapless) states around the Fermi surface even in the absence of disorder, which contradicts current experiments thus excluding such a pairing in iron pnictide superconductors.

¹This work was supported by the Texas Center for Superconductivity and the Robert A. Welch Foundation (Grant number E-1070) (Y.G. and W.P.S), and by U.S. DOE at LANL under Contract No. DE-AC52-06NA25396 (J.X.Z.)

Yi Gao
Department of Physics, University of Houston

Date submitted: 09 Nov 2009

Electronic form version 1.4