

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Superconducting fluctuations in one-dimensional quasi-periodic “metallic” chains. PAUL M. GRANT¹, W2AGZ Technologies — It is well known that a purely periodic chain of odd-electron atoms, nominally expected to exhibit metallic behavior, is unstable to charge/spin spatial displacement which lowers its ground state energy by gapping its multi-degenerate Fermi surface, in this case consisting of nesting parallel sheets. It is largely for these reasons that superconductivity is not observed in highly one-dimensional metals – it is simply energetically more favorable for CDW/SDW gaps to form, rather than a BCS state, at least one mediated by electron-phonon coupling. In this talk, we explore the hypothetical electronic properties of a nominally “metallic” quasi-periodic chain using both an analytical approach and computationally with density functional theory, searching for configurations which yield “gap-lets” sufficiently small to permit the formation of BCS pairs as the new energetically favored ground state. The particular embodiment we examine is a string of aluminum atoms with interatomic spacings determined by a Fibonacci sequence. We propose a path to attempt synthesis of such a structure for experimental examination. . . and perhaps leading to an entirely new class of higher temperature superconductors.

¹APS Senior Life Fellow

Paul M. Grant
W2AGZ Technologies

Date submitted: 18 Nov 2009

Electronic form version 1.4