

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Probing the symmetry of the pseudogap in cuprates superconductors with Angular Resolved Photoemission Spectroscopy (ARPES)¹

ARI D. PALCZEWSKI, TAKESHI KONDO, ADAM KAMINSKI, Ames Laboratory and Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA, J.S. WEN, G.Z.J. XU, G. GU, Condensed Matter Physics and Materials Science Department, Brookhaven National Laboratory, Upton, New York 11973, USA — There is still no consensus on the origins of the pseudogap in cuprate superconductors; recent studies have shown that it is probably not a preformed cooper pair state, but a completely different state that competes with superconductivity. Some of the possible different states include a charge density state where electrons scatter between a Q-vector connecting the nested bands close to $(\pi,0)$ or a spin density state that is derived from the (π,π) Q-vector. The difference between the two, or other exotic states, might be ascertained by looking at the symmetry of the pseudogap as a function of momentum space. We use Angular Resolved Photoemission Spectroscopy (ARPES) to probe this symmetry or lack thereof in cuprates.

¹Contract No. DE-AC02-07CH11358

Ari D. Palczewski
Ames Laboratory and Department of Physics and Astronomy,
Iowa State University, Ames, IA 50011, USA

Date submitted: 18 Nov 2009

Electronic form version 1.4