

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Role of Codeposited Impurities in Growth: Explaining Cu(0 0 1)¹

A. BH. HAMOUDA, Univ. of Maryland, College Park & Univ. of Monastir, Tunisia, RAJESH SATHIYANARAYANAN, UMD & Pennsylvania State Univ., University Park, A. PIMPINELLI, UMD & Science Attaché, French Embassy, Houston, T. L. EINSTEIN, UMD — A unified explanation of the physics underlying all the distinctive features of the growth instabilities observed on Cu vicinals² has long eluded theorists. Recently, kinetic Monte Carlo studies showed that codeposition of impurities during growth could account for all the experimental observations.³ To identify the responsible impurity atom, we compute the nearest-neighbor binding energies (E_{NN}) and terrace diffusion barriers (E_d) for several candidate impurity atoms on Cu(1 0 0) using DFT-based VASP. Our calculations show that codeposition (with Cu) of mid-transition elements, such as Fe, Mn, and W, could cause the observed instabilities; when the experimental set-up is considered, W emerges to be the most likely candidate. We discuss the role of impurities in nanostructuring of surfaces.

¹Supported by NSF MRSEC Grant DMR 05-20471; NSF supported computer usage at NCSA, UIUC.

²N. Néel . . . H.-J. Ernst., J. Phys.: Condensed Matter 15, S3227 (2003).

³A. Ben-Hammouda et al., Phys. Rev. B 77, 245430 (2008).

Rajesh Sathiyarayanan
Univ. of Maryland, College Park

Date submitted: 19 Nov 2009

Electronic form version 1.4