

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Point-contact spectroscopy of the iron chalcogenide superconductors: interplay between multiband superconductivity and magnetism¹
WAN KYU PARK, C.R. HUNT, H.Z. ARHAM, X. LU, L.H. GREENE, University of Illinois at Urbana-Champaign, Z.J. XU, J.S. WEN, Z.W. LIN, Q. LI, G. GU, Brookhaven National Laboratory — We report point-contact conductance measurements on the iron chalcogenide superconductors, $\text{Fe}_{1+y}\text{Te}_{1-x}\text{Se}_x$. The excess Fe atoms are known to occupy the interstitial sites in the Te-Se plane, affecting the superconductivity as well as the magnetism in this family. For a compound having nominal values of $y=0$ and $x=0.45$, a single superconducting transition is observed at 14.2 K. In the superconducting state, BTK-like double peak structures due to Andreev reflection are observed. However, the peak position of different point contacts falls to a wide voltage range, 1.5 – 4 mV. Additional multiple humps are sometimes observed in a much higher bias voltage range, 8 – 15 mV. Most strikingly, conductance enhancement persists well above T_c . We will present possible interpretations of these experimental observations in terms of multiband superconductivity and the interplay between superconductivity and magnetism.

¹The work at UIUC is supported by the U.S. DOE under Award No. DE-FG02-07ER46453 and by NSF-DMR-0706013 through FSMRL and CMM, and the work at BNL is carried out under U.S. DOE Award No. DE-AC0298CH10886.

Wan Kyu Park
University of Illinois at Urbana-Champaign

Date submitted: 24 Nov 2009

Electronic form version 1.4