

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Auto-correlation ARPES study of Pb-Bi2201 MAKOTO HASHIMOTO, Stanford University / LBL, RUIHUA HE, JEAN-PIERRE TESTAUD, WORAWAT MEEVASANA, ROB MOORE, DONGHUI LU, YOSHIYUKI YOSHIDA, HIROSHI EISAKI, THOMAS DEVEREAUX, ZAHID HUSSAIN, ZHIXUN SHEN — It is important to understand the electronic structure in momentum (k -) and real (r -) spaces in a unified picture. In high-temperature (T_c) cuprates, the different shape of superconducting gap anisotropies have been found by scanning tunneling microscopy (STM) and angle-resolved photoemission spectroscopy (ARPES). While the ARPES studies show simple d-wave superconducting gap around the node below T_c , the STM studies which approach superconducting gap anisotropy in k -space by employing the “octet model” show the deviation from simple d-wave superconducting gap especially in the underdoped region. To explore this issue, we study auto-correlation (AC-) ARPES spectra of optimally doped Pb-Bi2201 measured at $T \ll T_c$, $T_c < T \ll T^*$, and $T > T^*$. The obtained AC-ARPES spectra show very similar behavior to the reported Fourier transformed STM spectra. In the presentation, we will compare these ARPES and STM results and discuss the relationship between r - and k -space focusing on the superconducting gap anisotropy.

Makoto Hashimoto
Stanford University / LBL

Date submitted: 18 Dec 2009

Electronic form version 1.4