

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Time-dependent density-matrix functional theory formalism to study biexcitonic phenomena in bulk systems and nanostructures¹

VOLODYMYR TURKOWSKI, Dept. of Physics and NSTC, University of Central Florida, Orlando FL 32816, TALAT S. RAHMAN, Dept. of Physics and NSTC, University of Central Florida, Orlando, FL 32816, CARSTEN A. ULLRICH, Dept. of Physics and Astronomy, University of Missouri, Columbia, MO 65211, MICHAEL N. LEUENBERGER, Dept. of Physics and NSTC, University of Central Florida, Orlando, FL 32816 — We formulate a time-dependent density-matrix functional theory (TDDMFT) approach to study higher-order correlation effects like biexcitons in different bulk systems and nanostructures. In particular, we derive the TDDMFT version of the Schroedinger equation for biexcitons in terms of the one-body and two-body reduced density matrices. To test the approach, we calculate the biexcitonic binding energies in the case of different exchange- correlation (XC) potentials for bulk CuCl, CuBr, CdS and ZnO materials with rather large biexcitonic binding energies. We show that the excitonic, biexcitonic and other higher-order correlation effects are more pronounced in the case when the XC kernel contains a $1/q^2$ Coulomb singularity. Also, we analyze the role of non-adiabaticity of the XC potential in description of the higher-order correlation effects within the TDDMFT.

¹Work supported in part by DOE Grant No. DOE-DE-FG02-07ER15842 and NSF Grants No. ECCS 0725514 and No. ECCS-0901784.

Volodymyr Turkowski
Department of Physics and NSTC, University of Central Florida,
Orlando FL 32816

Date submitted: 01 Dec 2009

Electronic form version 1.4