

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Quadratic Electro-Optic Effect in Doped Nonconjugated Conductive Polymer 1,4-Trans-Polyisoprene, an Organic Quantum Dot System

S. SHRIVASTAVA, A. NARAYANAN, M. THAKUR, Photonic Materials Research Laboratory, Auburn University, AL 36849 — Thin, optically uniform films of 1,4-trans-polyisoprene have been prepared on quartz substrate from a toluene solution. These films have been characterized using FTIR and optical absorption spectroscopy before and after doping with iodine. The optical absorption spectrum at low doping shows two peaks: one at ~ 4.2 eV due to radical cation and the other at ~ 3.2 eV due to charge-transfer. Doping leads to a reduction of the intensity of =C-H bending vibration-band due to formation of radical cations upon charge-transfer. Quadratic electro-optic measurements have been made using field-induced birefringence method at 633 nm. A modulation depth of $\sim 0.13\%$ has been observed for an applied field of $1.1 \text{ V}/\mu\text{m}$ for a $0.37 \mu\text{m}$ thick film. The modulation depth had a quadratic dependence on applied field. The Kerr coefficient as measured is exceptionally large and has been attributed to the subnanometer size metallic domains (quantum dots) formed upon doping and charge-transfer.

M. Thakur
Photonic Materials Research Laboratory, Auburn University, AL 36849

Date submitted: 28 Nov 2009

Electronic form version 1.4